



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2008, 19th Annual JWP Conference

Apr 12th, 2:35 PM - 3:35 PM

The Role of Medial Septal Gabaergic Neurons in Prefrontal Cortex Theta Modulation

Ryan Peterson

Illinois Wesleyan University

Joseph Williams, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Ryan Peterson and Joseph Williams, Faculty Advisor, "The Role of Medial Septal Gabaergic Neurons in Prefrontal Cortex Theta Modulation" (April 12, 2008). *John Wesley Powell Student Research Conference*. Paper 20.
<http://digitalcommons.iwu.edu/jwprc/2008/posters2/20>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P38

**THE ROLE OF MEDIAL SEPTAL GABAERGIC NEURONS IN PREFRONTAL
CORTEX THETA MODULATION**

Ryan Peterson and Joseph Williams*

Psychology Department, Illinois Wesleyan University

Theta activity recorded in various structures of the brain including the hippocampus, amygdala and prefrontal cortex (PFC) has been associated with attention, memory, decision-making and emotions. While several researchers have determined that hippocampal theta is gated by medial septal area (MSA) neuronal firing, the same has not been equally demonstrated in the PFC. Previous research in this lab has shown MSA cholinergic neurons modulate PFC theta. However it is possible that other neurotransmitter systems, such as the MSA GABAergic system also help regulate PFC theta. Given that hippocampal theta is gated by both acetylcholine and GABA, the GABAergic connections from the MSA to the PFC must be determined. Eight male Long Evans rats were used in the present experiments. A guide cannula was surgically placed in the MSA and a recording electrode was placed in the PFC. Several days later the rats were anesthetized with an injection of ketamine/xylazine and then infused with various doses of ethanol and the GABAergic agonist muscimol. The researchers predicted that both ethanol and muscimol in the MSA will reduce or eliminate theta activity in the PFC, similar to their effects on hippocampal theta.